

Relations for Direct CP asymmetries in $B \rightarrow PP$ and $B \rightarrow PV$ decays¹

T. N. Pham

*Centre de Physique Théorique,
Centre National de la Recherche Scientifique, UMR 7644,
Ecole Polytechnique, 91128 Palaiseau Cedex, France*

ABSTRACT

The presence of additional strong phase from power corrections and other chirally enhanced terms makes it more difficult to predict direct CP asymmetries in two-body charmless B decays. In this talk, I would like to report on a recent work on QCD Factorisation and Power Corrections in Charmless B Decays. Using the measured branching ratios for $B \rightarrow PV$, it is shown that power corrections in charmless B decays are probably large, at least for penguin dominated PV channels. Since the tree-penguin interference responsible for direct CP asymmetries in two-body charmless B decays are related by CKM factors and $SU(3)$ symmetry, we find that, if power corrections other than the chirally enhanced power corrections and annihilation topology were negligible, QCD Factorisation would predict the direct CP asymmetry of $B \rightarrow \pi^+\pi^-$ to be about 3 times larger than that of $B \rightarrow \pi^\pm K^\mp$, with opposite sign, in agreement with the latest measurement from Belle. Similar relations are also given for direct CP asymmetries in $B \rightarrow PV$.

¹Talk given at the QCD@Work 2005, Conversano, Bari, Italy, 16-20 June 2005

1 Introduction

The large direct CP asymmetries observed at BaBar and Belle in $B \rightarrow K\pi$ and $B \rightarrow \pi\pi$ decays [1, 2] indicate a large strong phase in these decay amplitudes. Since in general QCD Factorisation(QCDF) predicts a much smaller strong phase and a small CP asymmetry, one would then need important power correction terms or other power-suppressed non-factorisable term (e.g FSI effects etc.) to generate a large strong phase and a large CP asymmetry in these decays. In fact the charmed meson inelastic rescattering FSI effects[3, 4] or charming penguin[5] are able to produce a large absorptive part and therefore a large strong phase for $B \rightarrow PP$ and $B \rightarrow PV$ amplitudes. Though the presence of these power-suppressed terms makes it difficult to predict the amount of CP asymmetry, because of $SU(3)$ symmetry and the CKM factor, one can derive however relations between CP asymmetries in $B \rightarrow PP$ and $B \rightarrow PV$. In the following I shall first present an analysis showing possible evidence for power-suppressed terms in charmless B decays.

In QCD Factorization (QCDF)[6], the $O(1/m_b)$ power corrections in penguin matrix elements and other chirally enhanced corrections could make important contributions to the penguin-dominated charmless B decays as in $B \rightarrow \pi K$ decays. Other power corrections terms such as annihilation contributions may also be present in PP and PV decays as first noticed in the perturbative QCD method for charmless B decays[7] and indicated by recent analysis of charmless two-body non-leptonic B decays[8, 9, 10]. In a recent work[11], we have shown that in QCDF, it is possible to consider certain ratios of the $B \rightarrow PV$ amplitudes which depend only on the Wilson coefficients and the known hadronic parameters. The discrepancy between prediction and experiment for the ratio would be a clear evidence for annihilation or other non-factorisable contributions. We find that annihilation topology likely plays an indispensable role at least for penguin-dominated PV channels. Including the annihilation terms in QCDF, we find that the direct CP asymmetry of $B \rightarrow \pi^+\pi^-$ to be about 3 times larger than that of $B \rightarrow K^\mp\pi^\pm$, with opposite sign, in agreement with experiment[2].

2 QCD factorization for charmless B decays

The effective Lagrangian for non-leptonic B decays can be obtained from operator product expansion and renormalization group equation, in which short-distance effects involving large virtual momenta of the loop corrections from the scale M_W down to $\mu = \mathcal{O}(m_b)$ are integrated into the Wilson coefficients. The amplitude for the decay $B \rightarrow M_1 M_2$ can be expressed as

$$\mathcal{A}(B \rightarrow M_1 M_2) = \frac{G_F}{\sqrt{2}} \sum_{i=1}^6 \sum_{q=u,c} \lambda_q C_i(\mu) \langle M_1 M_2 | O_i(\mu) | B \rangle \quad (1)$$

λ_q is a CKM factor, $C_i(\mu)$ are the Wilson coefficients perturbatively calculable from first principles and O_i are the tree and penguin operators given by(neglecting other operators):

$$\begin{aligned} O_1 &= (\bar{s}u)_L(\bar{u}b)_L \quad , \quad O_4 = \sum_q (\bar{s}q)_L(\bar{q}b)_L \\ O_6 &= -2 \sum_q (\bar{s}_L q_R)(\bar{q}_R b_L) \end{aligned} \quad (2)$$

The hadronic matrix elements $\langle M_1 M_2 | O_i(\mu) | B \rangle$ contain the physics effects from the scale $\mu = \mathcal{O}(m_b)$ down to Λ_{QCD} . In the heavy quark limit, QCD Factorisation [6] allows the decay mplitude

$\langle M_1 M_2 | O_i(\mu) | B \rangle$ to be factorized into hard radiative corrections and non-perturbative matrix elements which can be parametrized by the semi-leptonic decays form factors and meson light-cone distribution amplitudes (LCDAs).

Power corrections in $1/m_b$ come from penguin matrix elements, chirally enhanced corrections and annihilation contributions. For example, in the $B \rightarrow \pi K$ amplitude, the matrix element of O_6 is of the order $O(1/m_b)$ compared to the $(V-A) \times (V-A)$ O_1 and O_4 matrix elements, since $\langle K | \bar{s}_L d_R | 0 \rangle$ is proportional to $m_K^2/m_s \approx 2.5 \text{ GeV}$ while $\langle K | \bar{s}_L d_L | 0 \rangle$ is proportional to K momentum which is $O(m_b)$, thus numerically, the matrix element of O_6 has a factor

$$r_\chi^K = \frac{2m_K^2}{m_b(m_s + m_d)} \approx O(1) \quad (3)$$

and is comparable to that of O_4 . For penguin-dominated decays, the O_4 and O_6 matrix element are of the same sign in PP channel, while in PV channel they are of opposite sign. Thus in QCDF one expects a small $B \rightarrow K\rho$ branching ratio relative to $B \rightarrow \pi K$. Because of a cancellation between the O_4 and O_6 contributions, the $B \rightarrow K\rho$ decay is more sensitive to other power corrections and non-factorisable contributions. Including the chirally-enhanced corrections in terms of two quantities $X_{A,H}$ and a strong phase, the $B \rightarrow M_1 M_2$ decay amplitudes in QCDF can be thus be written as[12, 13]:

$$\mathcal{A}(B \rightarrow M_1 M_2) = \frac{G_F}{\sqrt{2}} \sum_{p=u,c} V_{pb} V_{ps}^* \left(- \sum_{i=1}^6 a_i^p \langle M_1 M_2 | O_i | B \rangle_f + \sum_j f_B f_{M_1} f_{M_2} b_j \right), \quad (4)$$

3 Power corrections in $B \rightarrow PV$ decays

Consider the ratio of $A(B^+ \rightarrow \pi^+ K^{*0})$ to $A(B^0 \rightarrow \rho^+ \pi^-)$. amplitudes. If the power corrections were negligible, this ratio would be theoretically very clean where the form factors cancel out, furthermore it is almost independent of the CKM angle γ and the strange-quark mass:

$$\left| \frac{\mathcal{A}(B^+ \rightarrow \pi^+ K^{*0})}{\mathcal{A}(B^0 \rightarrow \rho^+ \pi^-)} \right| \simeq \left| \frac{V_{cb} V_{cs}}{V_{ub} V_{ud}} \right| \frac{f_{K^*}}{f_\rho} \left| \frac{a_4^c(\pi K^*) + r_\chi^{K^*} a_6^c(\pi K^*)}{a_1^u} \right| \quad (5)$$

$|(a_4^c(\pi K^*) + r_\chi^{K^*} a_6^c(\pi K^*))/a_1^u|$ should be about or less than 0.04 in QCDF. ($f_{K^*}/f_\rho \approx 1$). The ratio $|V_{ub}/V_{cb}|$ is not very well determined experimentally, but a stringent lower limit can be obtained from the unitarity of the CKM matrix . Since [14, 15] :

$$\left| \frac{V_{ub}}{V_{cb}} \right| = \lambda \sin \beta \sqrt{1 + \frac{\cos^2 \alpha}{\sin^2 \alpha}} \geq \lambda \sin \beta. \quad (6)$$

and from the current Babar and Belle measured values : $\sin 2\beta = 0.725 \pm 0.037$ [16] , we have

$$\left| \frac{V_{ub}}{V_{cb}} \right| \geq \lambda \sin \beta = 0.090 \pm 0.007 > 0.078 \quad (7)$$

Eq.(5) implies the following inequality :

$$0.53 > \left| \frac{\mathcal{A}(B^+ \rightarrow \pi^+ K^{*0})}{\mathcal{A}(B^0 \rightarrow \rho^+ \pi^-)} \right| = 0.77 \pm 0.09, \quad (8)$$




Figure 1: The ratio $\mathcal{B}(B^+ \rightarrow \pi^+ K^{*0})/\mathcal{B}(B^0 \rightarrow \rho^+ \pi^-)$ versus the weak annihilation phase ϕ_A . The default parameters are used but letting the annihilation parameter $\rho_A = 1$. The dashed lines show the ratios without weak annihilation contributions. The gray areas denote the experimental measurements with 1σ error.




Figure 2: The ratio $\mathcal{B}(B^0 \rightarrow K^+ \rho^-)/\mathcal{B}(B^0 \rightarrow \rho^- \pi^+)$ versus the weak annihilation phase ϕ_A . The default parameters are as in Fig.1

where the number on the rhs is from the measured branching ratios [17, 18]. The lhs would be reduced further to 0.46 ± 0.04 , if in Eq.(6) one neglects a small $\cos^2 \alpha$ term according to a recent determined value $\alpha = (101^{+16}_{-9})^\circ$ [16].

Since the chirally enhanced corrections for penguin-dominated decays are not expected to be large, this large discrepancy is strong indication that annihilation topology and/or other sources of power corrections might play an important role at least in $B \rightarrow PV$ decays. There is similar disagreement between theory and experiment in another ratio, the branching fraction of $B^0 \rightarrow K^+ \rho^-$ to that of $B^0 \rightarrow \rho^- \pi^+$, though with large theoretical uncertainties. For $\gamma = 70^\circ$, $V_{ub}/V_{cb} = 0.09$, $a_4^c(\rho K) - r_\chi^K a_6^c(\rho K) = 0.037 + 0.003i$, $m_s = 90$ MeV, we find

$$\frac{\mathcal{B}(B^0 \rightarrow K^+ \rho^-)}{\mathcal{B}(B^0 \rightarrow \rho^- \pi^+)} = 0.38 \quad (9)$$

far below the measured value of 1.01 ± 0.34 , though, this ratio could be increased to 0.69, if m_s is lowered to 70 MeV. The discrepancy could be greater according to a recent Belle measurements[17] which give a very large $B^0 \rightarrow K^+ \rho^-$ branching ratio, $(15.1^{+3.4+2.4}_{-1.5-2.6}) \times 10^{-6}$

which could be obtained by a large annihilation contributions[13]. Recent charmed meson in-elastic rescattering FSI calculation[4] also produces a large branching ratio. Taken together, these results indicate that the penguin-dominated $B \rightarrow PV$ decay amplitudes are consistently underestimated without annihilation contributions. Including the annihilation terms, from Eq. (4), we have

$$\begin{aligned} A(B^+ \rightarrow \pi^+ K^{*0}) &= f_{K^*} F^{B\pi} m_B^2 a_4 + b_3(V, P) \\ A(B^0 \rightarrow K^+ \rho^-) &= f_K A_0^{B\rho} m_B^2 (a_4 - r_\chi^K a_6) + b_3(P, V) \end{aligned} \quad (10)$$

$$b_3(M_1, M_2) = \frac{C_F}{N_c^2} \{C_3 A_1^i(M_1, M_2) + C_5 A_3^i(M_1, M_2) + (C_5 + N_c C_6) A_3^f(M_1, M_2)\} \quad (11)$$

With the penguin terms $a_4 \simeq -0.03$ and $a_4 - r_\chi^K a_6 \simeq 0.037$ having opposite sign, the key observation is that $b_3(V, P)$ and $b_3(P, V)$, which get most of the contribution from $(C_5 + N_c C_6) A_3^f$ term, are also roughly of the opposite sign since $A_3^f(P, V) = -A_3^f(V, P)$. Thus QCDF can easily enhance both ratios without fine tuning (no large strong phase) as can be seen in Fig.2 .

4 Direct CP violations

We now turn to the CP asymmetries in QCDF with annihilation terms included. Because of the CKM factor and $SU(3)$ symmetry for the tree and penguin matrix elements in $B^0 \rightarrow \pi^+ \pi^-$ and $B^0 \rightarrow K^+ \pi^-$ decays, one can derive a relation between direct CP asymmetries in these two channels. With the CP asymmetry given as:

$$\begin{aligned} A_{\pi\pi} &= \frac{4|V_{ub}V_{ud}V_{cb}V_{cd}T_{\pi\pi}P_{\pi\pi}|\sin\gamma\sin\delta}{2\mathcal{B}(B \rightarrow \pi^+\pi^-)}, \\ A_{\pi K} &= -\frac{4|V_{ub}V_{us}V_{cb}V_{cs}T_{\pi K}P_{\pi K}|\sin\gamma\sin\tilde{\delta}}{2\mathcal{B}(B \rightarrow \pi^+K^-)}. \end{aligned} \quad (12)$$

($\delta = \delta_P - \delta_T$ = strong phases difference between the penguin and tree amplitudes), we find

$$\begin{aligned} \frac{A_{\pi\pi}}{A_{\pi K}} &= -\frac{f_\pi^2}{f_K^2} \frac{\mathcal{B}(B \rightarrow \pi^+ K^-)}{\mathcal{B}(B \rightarrow \pi^+ \pi^-)} \left| \frac{T_{\pi\pi} P_{\pi\pi}}{T_{\pi K} P_{\pi K}} \right| \frac{\sin\delta}{\sin\tilde{\delta}} \\ &\simeq (-2.7 \pm 0.3) \frac{\sin\delta}{\sin\tilde{\delta}} \end{aligned} \quad (13)$$

a consequence of the fact that $T_{\pi\pi}P_{\pi\pi}/T_{\pi K}P_{\pi K}$ is close to 1, a reasonable approximation in QCDF, at about 10 percent level uncertainty. A previous derivation of this relation is given in [20, 21]. Belle has claimed large direct CP asymmetry observed in $B^0 \rightarrow \pi^+ \pi^-$ decay while BaBar has not confirmed it yet, but both of them are close in measurements on $A_{CP}(\pi^- K^+)$ [2, 1, 19]

$$A_{\pi\pi} = \begin{cases} 0.56 \pm 0.12 \pm 0.06 & (\text{Belle}) , \\ 0.09 \pm 0.15 \pm 0.04 & (\text{BaBar}) , \\ 0.37 \pm 0.10 & (\text{Average}) . \end{cases} \quad (14)$$

$$A_{\pi K} = \begin{cases} -0.101 \pm 0.025 \pm 0.005 & (\text{Belle}) , \\ -0.133 \pm 0.030 \pm 0.009 & (\text{BaBar}) , \\ -0.114 \pm 0.020 & (\text{Average}) . \end{cases} \quad (15)$$

We thus expect very naturally a larger direct CP violation for $\pi^+\pi^-$ decay compared with π^-K^+ decay, since the $\pi^+\pi^-$ decay rate is smaller than the π^-K^+ decay rate by factor 3 – 4,

Experimentally,

$$\frac{A_{\pi\pi}}{A_{\pi K}} = \frac{0.37 \pm 0.10}{-0.11 \pm 0.03} = -3.4 \pm 1.5 , \quad (16)$$

still consistent with the theoretical estimation of -2.7 ± 0.3 .

Similar relation between CP asymmetries for the $B \rightarrow PV$ decays for which the CP-violating interference terms are essentially of the same magnitude, but with opposite sign:

$$\begin{aligned} \frac{A_{\text{CP}}(B^0 \rightarrow \rho^+ \pi^-)}{A_{\text{CP}}(B^0 \rightarrow K^{*+} \pi^-)} &\simeq - \frac{\mathcal{B}(B^0 \rightarrow K^{*+} \pi^-)}{\mathcal{B}(B^0 \rightarrow \rho^+ \pi^-)} \frac{f_\rho^2}{f_{K^*}^2} \frac{\sin \delta_{\pi\rho}}{\sin \delta_{\pi K^*}} \\ \frac{A_{\text{CP}}(B^0 \rightarrow \rho^- \pi^+)}{A_{\text{CP}}(B^0 \rightarrow \rho^- K^+)} &\simeq - \frac{\mathcal{B}(B^0 \rightarrow \rho^- K^+)}{\mathcal{B}(B^0 \rightarrow \rho^- \pi^+)} \frac{f_\pi^2}{f_K^2} \frac{\sin \delta_{\rho\pi}}{\sin \delta_{\rho K}} \end{aligned} \quad (17)$$

In the presence of charming penguin or charmed meson inelastic rescattering FSI effects, the above CP asymmetry relation applies since the tree-penguin interference terms are related by $SU(3)$ symmetry and CKM factor[3]. Thus any significant deviation from theoretical estimation would suggest either different strong phases, e.g between $\pi\pi$ and πK decays, or possible new physics contributions.

5 Conclusion

Power corrections in charmless B decays are probably large, at least for the penguin-dominated PV channel. The key observation is that QCDF predicts the annihilation terms for $B^+ \rightarrow \pi^+ K^{*0}$ and $B^0 \rightarrow K^+ \rho^-$ to be almost equal in magnitude but opposite in sign and thus enhance the decay rates for these two modes to accommodate the experimental data. The relation for the direct CP asymmetry would naturally implies a large CP asymmetry for $B \rightarrow \pi^+\pi^-$, about 3 times larger than that of $B \rightarrow \pi^\pm K^\mp$ with opposite sign.

I would like to thank G. Nardulli, P. Colangelo, F. De Fazio and the organisers of QCD@Work for the warm hospitality extended to me at Conversano.

References

- [1] BaBar Collaboration, B. Aubert *et al.*, hep-ex/0501071.
- [2] Belle Collaboration, K. Abe *et al.*, hep-ex/0502035.
- [3] C. Isola, M. Ladisa, G. Nardulli, T. N. Pham and P. Santorelli, Phys. Rev. D **64**, 014029 (2001); *ibid*, **65**, 094005 (2002).
- [4] C. Isola, M. Ladisa, G. Nardulli, and P. Santorelli, Phys. Rev. D **68**, 114001 (2003) .
- [5] M. Ciuchini, E. Franco, G. Martinelli, and L. Silvestrini, Phys. Lett. **B 515**, 33 (2001) ; and previous works cited therein.

- [6] M. Beneke, G. Buchalla, M. Neubert and C. T. Sachrajda, Phys. Rev. Lett. **83**, 1914 (1999); Nucl. Phys. B **591**, 313 (2000) .
- [7] Y. Y. Keum, H. N. Li and A. I. Sanda, Phys. Rev. D **63**, 054008 (2001) ; Phys. Lett. B **504**, 6 (2001) .
- [8] D. S. Du, J. F. Sun, D.S. Yang and G. H. Zhu, Phys. Rev. D **67**, 014023 (2003) .
- [9] R. Aleksan, P. F. Giraud, V. Morenas, O. Pene and A. S. Safir, Phys. Rev. D **67**, 094019 (2003) .
- [10] N. de Groot, W. N. Cottingham and I. B. Whittingham, Phys. Rev. D **68**, 113005 (2003).
- [11] T. N. Pham and Guohuai Zhu, Phys. Rev. D **69** , 114016 (2004).
- [12] M. Beneke, G. Buchalla, M. Neubert and C. T. Sachrajda, Nucl. Phys. B **606**, 245 (2001).
- [13] M. Beneke and M. Neubert, Nucl. Phys. B **675**, 333 (2003) .
- [14] T. N. Pham, invited talk at the 2nd Workshop on the CKM Unitarity Triangle, Durham, April 2003, hep-ph/0306271 .
- [15] A. J. Buras, F. Parodi and A. Stocchi, JHEP **0301**, 029 (2003) .
- [16] J. Smith, Talk given at the 3rd Workshop on the CKM Unitarity Triangle, San Diego, USA, March 2005 .
- [17] Heavy Flavor Averaging Group(HFAG), hep-ph/0505100 .
- [18] Heavy Flavor Averaging Group(HFAG),
www.slac.stanford.edu/xorg/hfag/triangle/ichep2004/index.shtml .
- [19] Heavy Flavor Averaging Group(HFAG),
www.slac.stanford.edu/xorg/hfag/triangle/summer2005/index.shtml .
- [20] R. Fleischer, Phys. Lett. B **459** (1999) 306 ; A. J. Buras, R. Fleischer, S. Recksiegel and F. Schwab, hep-ph/0402112 .
- [21] M. Dariescu, N. G. Deshpande, X. G. He, and G. Valencia, Phys. Lett. B **557**, 60 (2003).